

Land-Based Solutions for Plastics in the Sea

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 101003954

D6.5 Environmental Risk Assessment (E.R.A.) of plastics
in aquatic and terrestrial environments

Due date of deliverable: 31/08/2024

Revised due date: 01/12/2024

Actual submission date: 29/11/2024



Horizon 2020
European Union Funding
for Research & Innovation

PROJECT INFORMATION

Project number: 101003954

Project acronym: LABPLAS

Project full title: Land-Based Solutions for Plastics in the Sea

Call: H2020-SC5-2018-2019-2020 submitted for H2020-SC5-2020-2 / 03 Sep 2020

Topic: CE-SC5-30-2020 – Plastics in the environment: understanding the sources, transport, distribution and impacts of plastics pollution

Type of action: RIA – Research and Innovation Action

Starting date: June 1st, 2021

Duration: 48 months

List of participants:

Nº	Participant name	Acronym	Country	Type
1	UNIVERSIDADE DE VIGO	UVI	SPAIN	HES
2	UNIVERSIDADE DA CORUÑA	UDC	SPAIN	HES
3	Bundesanstalt fuer Gewaesserkunde	BfG	GERMANY	RTO
4	LABORATORIO IBERICO INTERNACIONAL DE NANOTECNOLOGIA	INL	PORTUGAL	RTO
5	KATHOLIEKE UNIVERSITEIT LEUVEN	KUL	BELGIUM	HES
6	HELMHOLTZ ZENTRUM FUR OZEANFORSCHUNG KIEL	GEOMAR	GERMANY	RTO
7	NATIONAL OCEANOGRAPHY CENTRE	NOC	UNITED KINGDOM	RTO
8	SORBONNE UNIVERSITE	SU	FRANCE	HES
9	OPEN UNIVERSITEIT NEDERLAND	OUNL	NETHERLANDS	HES
10	LEIBNIZ INSTITUTE FOR BALTIC SEA RESEARCH	IOW	GERMANY	RTO
11	ASSOCIACAO PARA O DESENVOLVIMENTO DO ATLANTIC INTERNATIONAL RESEARCH CENTRE	AC	PORTUGAL	RTO
13	BASF SE	BASF	GERMANY	LE
14	TG ENVIRONMENTAL RESEARCH	ER	UNITED KINGDOM	SME
15	CONTACTICA S.L.	CTA	SPAIN	SME
16	STICHTING EGI	EGI	NETHERLANDS	Non-P
17	STICHTING RADBOUD UNIVERSITEIT	RU	NETHERLANDS	HES
18	UNIVERSIDADE FEDERAL DO PARÁ	UFPA	BRAZIL	HES



















DELIVERABLE DETAILS

Document number:	D6.5
Document title:	Environmental Risk Assessment (E.R.A.) of plastics in aquatic and terrestrial environments
Dissemination level	PU – Public
Period:	RP3
WP:	WP6
Task:	Task 6.4
Status:	Final Report
Coordinator:	R. Beiras
Other authors:	B. Espiña, C. Scherer, J. Dominguez, P. Baudy-Groh, A. Vilas, K. Voges
Recommended citation format	R. Beiras, B. Espiña, C. Scherer, J. Dominguez, P. Baudy-Groh, A. Vilas, K. Voges, 2024, Environmental Risk Assessment of plastics in aquatic and terrestrial environments, Deliverable 6.5, LABPLAS Grant Agreement No. 101003954 H2020-SC5-2020-2
Executive summary:	<p>According to the standard methodology, environmental risk posed by microplastics was quantitatively characterized using the following expression: Risk Quotient (R) = Exposure Level (EL) / Toxicity Threshold (TT). With that aim, the microplastic concentrations measured in the LABPLAS Project Work Package (WP) 2 in freshwater and marine aquatic environmental compartments were compared with the ecotoxicity thresholds found in the laboratory toxicity tests performed in WP6 using organisms representative of different trophic levels for each environmental compartment. A deterministic approach using an Assessment Factor (AF) and the toxicity threshold of the most sensitive test species, and a probabilistic approach based on the Species Sensitivity Distribution (SSD) curve were both conducted. Even under very conservative assumptions, such as using data from hotspots, AF=1000, toxicity thresholds obtained from most sensitive species, or taking the lower end of the 95% confidence interval (CI), R values were always <1; from 0.004 to 0.100 (deterministic approach) or 0.079 (probabilistic approach). Therefore, levels of risk in both freshwater and marine habitats are not concerning. Although the LABPLAS Project did not sample terrestrial microplastics, a preliminary attempt to quantify risk in soils was made by obtaining exposure levels from published data. In contrast to aquatic results, for terrestrial agriculture</p>

soils R values obtained with the maximum levels reported in the literature **were >1**. This is partly due to the higher sensitivity of *Eisenia sp.* (earthworm) compared to aquatic model species, but mainly due to the much higher plastic loads reported for some agriculture soils compared to aquatic environments.

Version	Date	Comments
1.0	07.03.2024	First draft
2.0	13.11.2024	Evaluation draft for partners
3.0	29.11.2024	Final version

Disclaimer

The views and opinions expressed in this document reflect only the authors' views, and not necessarily those of the European Commission.

TABLE OF CONTENTS

PROJECT INFORMATION	1
DELIVERABLE DETAILS	2
TABLE OF CONTENTS.....	4
LIST OF FIGURES AND TABLES	5
ABBREVIATIONS AND ACRONYMS.....	6
1 INTRODUCTION	7
2 METHODOLOGY	7
2.1 Exposure level assessment.....	7
2.2 Toxicity threshold assessment.....	8
2.3 Characterization of Risk	9
3 RESULTS	9
3.1 Exposure levels.....	9
3.1.1 Exposure levels in the river water column.....	9
3.1.2 Exposure levels in the North Sea surface water	10
3.1.3 Exposure levels in the river sediments.....	10
3.1.4 Estimation of mass using actual weight records	11
3.1.5 Exposure levels in agricultural soils	11
3.2 Toxicity thresholds.....	11
3.2.1 Aquatic species.....	13
3.2.2 Terrestrial species	13
3.3 Deterministic estimation of risk	13
3.3.1 Freshwater sites	14
3.3.2 Marine sites.....	14
3.3.3 Sediments.....	14
3.3.4 Soil	15
3.4 Probabilistic Estimation of Risk.....	15
4 DISCUSSION	16

LIST OF FIGURES

Figure 1. LABPLAS Project plastic toxicity test battery applied to test environmental field samples.	8
Figure 2. SSD curves for the most toxic plastic samples in aquatic (ID124) and terrestrial (ID131) habitats respectively.	15

LIST OF TABLES

Table 1. Abundance of floating plastic particles between 10 µm and 1 mm in the water of the Thames and Elbe rivers.....	10
Table 2. Abundance of floating plastic particles between 10 µm and 1 mm in the water of the North Sea	10
Table 3. Abundance of plastic particles between 10 µm and 1 mm in the sediments of the Thames and Elbe rivers.....	11
Table 4. Summary of the toxicity tests conducted with micronized environmental plastics sampled in the different areas studied in the LABPLAS Project in Germany (GE), Spain (ES) and the United Kingdom (UK). Test species and endpoints included <i>Vibrio fischeri</i> bioluminescence, sea-urchin embryo test (SET), freshwater and marine cyanobacteria/algae population growth (<i>Microcystis aeruginosa</i> and <i>Skeletonema</i> sp. respectively), freshwater and marine invertebrate reproduction and larval survival (<i>Daphnia magna</i> and <i>Acartia</i> sp respectively), development of early life stages of freshwater and marine fish (<i>Danio rerio</i> and <i>Cyprinodon variegatus</i> respectively), soil microbiota respiration, plant seedling emergence (<i>Lepidium sativum</i>) and earthworm reproduction (<i>Eisenia andrei</i>). L: leachate test; P: particle test. Results have been arbitrarily classified into categories of increasing toxicity indicated by green, yellow, orange and red. Methods have been thoroughly described in previous deliverables from D6.1 to D6.4.....	12
Table 5. Toxicity exerted by the most toxic freshwater plastics (ID-124) on the microscale and habitat-specific tests. L: leachate test. All leachates were made up at 10 g/L except for <i>Daphnia</i> (1 g/L).....	13
Table 6. Toxicity exerted by the most toxic terrestrial plastics (ID-131) on the microscale and terrestrial tests. L: leachate test; P: particle test.....	13
Table 7. Risk Quotient (R) obtained according to the deterministic method for the freshwater sites.....	14
Table 8. Risk Quotient (R) obtained according to the deterministic method for the marine sites.....	14

ABBREVIATIONS AND ACRONYMS

Abbreviation / Acronym	Description
AF	Assessment Factor
C	Concentration
CI	Confidence Interval
D	Deliverable
EC	European Commission
EC₁₀	Effect Concentration at which a 10% effect is observed
EL	Exposure Level
EPS	Extracellular Polymeric Substances
ERA	Environmental Risk Assessment
HC₅	Hazardous Concentration for 5% of species
LOEC	Lowest Observed Effect Concentration
MP	Microplastics
NOEC	No Observed Effect Concentration
PS	Poly-Styrene
R	Risk Quotient
SET	Sea urchin Embryo Test
SMP	Small Microplastics (10 to 1000 µm)
SMNP	Small Micro- and Nano- Plastics
spp	Species
std	Standard Deviation
SSD	Species Sensitivity Distribution
TT	Toxicity Threshold
WP	Work Package

1 INTRODUCTION

Despite the numerous benefits plastics provide to society, valid concerns exist regarding the harmful effects posed by their fragmentation into smaller particles. The smaller the size the higher the risk posed by these particles to organisms and human health. Small micro- and nano- plastics (SMNP) have been found across all environmental compartments, including air, freshwater, marine environments, and soils. For this reason, this document aims to perform a comprehensive ecological risk assessment (ERA) of SMNPs across various environmental compartments.

With this aim, two approaches will be followed. First, according to a mechanistic approach, risk can be assessed using **Risk Quotients** (R), which can be calculated by comparing microplastic (MP) densities in different environmental compartments, as obtained from Work Package (WP) 2 (refer to Deliverables D2.2, D2.3, D2.4, and unpublished data from 2023 cruises) with toxicity thresholds (TT) derived from laboratory tests using organisms representative of different trophic levels in each relevant environmental compartment, as conducted by WP6 (refer to D6.3 & D6.4). Since terrestrial habitats were not sampled for SMNPs in the LABPLAS project, exposure levels will be estimated using data from the literature.

Secondly, a probabilistic approach will be used to compare the distribution of MP abundance detected in the different environmental compartments sampled by WP2 with the Species Sensitivity Distribution (SSD) derived from WP6 laboratory tests for aquatic and terrestrial organisms.

Furthermore, this document refines previous ERA studies by incorporating ecotoxicological data derived from environmental plastics, used to generate secondary microplastics of heterogeneous composition, size and morphology, representative of actual environmental MPs, rather than by resorting to engineered microbeads as most of the earlier studies.

2 METHODOLOGY

The ERA was conducted following the standard methodology established by the United States National Academy of Sciences¹, initially developed for Human Health Risk Assessment to address threats to human health caused by occupational exposure. Risk is quantitatively characterized as a Risk Quotient (R) comparing two estimates: **Exposure Level** (EL) to **Toxicity Thresholds** (TT). Thus, $R = EL / TT$. The closer the exposure levels are to the toxicity thresholds, the higher the associated risk, with values of $R > 1$ not acceptable.

2.1 Exposure level assessment.

Exposure levels for aquatic habitats were estimated based on the measured SMP (>100 µm) concentrations recorded by WP2 in each sampled environmental compartment. Filtered particles were analyzed by the LABPLAS project partner NOC using the methods described in D2.1 and D2.4. Mass data were derived from particle counts and bi-dimensional data using two approaches: the estimates provided by the siMPle software (<https://simple-plastics.eu/>), and those obtained from the actual measurements of mass vs area in plastic particles, down to a maximum dimension of 0.4 mm, conducted by GEOMAR.

Since the LABPLAS Project does not target sampling microplastics in terrestrial habitats, EL for soils was obtained from the scientific literature.

¹ NRC 1983. Risk assessment in the Federal Government: managing the process. National Research Council. National Academy Press, Washington DC.

The contents of this document are the copyright of the LABPLAS consortium and shall not be copied in whole, in part, or otherwise reproduced, used, or disclosed to any other third parties without prior written authorisation.

2.2 Toxicity threshold assessment

Small, micro and nano plastics (SMNPs) were produced from field plastic samples obtained from three different environmental compartments: marine (shoreline and sea surface), freshwater (river-banks and surface water), and terrestrial (soil), according to methods described in D6.1. and D6.3. To identify the hazard and assess the risk posed by these SMNPs, a battery of aquatic and terrestrial ecotoxicological tests (Fig. 1), selected according to internationally accepted standardised methods adapted to plastic toxicity testing, and using organisms representative of different trophic levels and environmental compartments, were conducted.




Figure 1. LABPLAS Project plastic toxicity test battery applied to test environmental field samples.

As previously described, the LABPLAS plastic toxicity assessment scheme takes into account two routes of exposure: (1) waterborne chemicals leached from the plastics (Tier I), and (2) plastic particles in contact and/or ingested by the organisms (Tier II) (Figure 3, D6.4). Tier I exposes the organisms to serial dilutions of a leachate obtained by following the standard protocol by Almeda et al, (2023), using a load of 10 g of plastic per liter. Leaching time is standardized to 24 h consistently with Dhavamani et al. (2022) who found maximum concentrations of leached plastic additives after 24-48 h followed by a decrease due to readsorption. Leachate testing allows the detection of effects caused by chemicals leached from the plastics into the water. These chemicals may be part of the original plastic composition or, in the case of environmental plastics, sorbed from the surrounding aquatic medium. Tier II exposes the organisms to plastic particles of a standard size suitable to be ingested by the test species and natural particles of an analogous size range. This size depends on the model organism. For most filter feeders and early stages of aquatic organisms, an upper size limit of 20 μm is suited.

Depending on the completeness of the data sets, TT were estimated according to the following statistical methods. If the data set allowed fitting to a concentration:response non-linear model, the TT was estimated as the EC_{10} , or concentration of MPs reducing by 10% the biological endpoint recorded. In other cases, treatments

were statistically compared to the control and the lowest observed effect concentration (LOEC) was taken as TT estimation.

For Tier I tests, dilutions were translated into mass concentration units by multiplying the dilution factor by the concentration used to produce the tested leachates; either 1 or 10 g/L.

2.3 Characterization of Risk

Risk was characterized using both a deterministic approach, based on the TT of the most sensitive test species of each habitat (critical value), and a probabilistic approach based on the Species Sensitivity Distribution (SSD) curve.

In the deterministic approach,

$$R = EL \times AF / TT$$

Where AF is an assessment factor (AF) – also called uncertainty factor – intended to make conservative allowances for the various sources of uncertainty associated with both EL and TT estimation. Typical sources of uncertainty are extrapolation of chronic effects from acute toxicity data, laboratory-to-field extrapolations, or allowances for the representativeness of the chosen test species. Typical values of AF range from 10 to 1000, decreasing as the amount and quality of ecotoxicological information increases. The [EC](#) prescribes a value of 10 when long-term effects in three trophic levels are assessed, increasing to a value of 1000 when only TT for acute effects are available. Obviously, high AF values increase the risk of too conservative – and thus, in practice, hardly applicable – estimates of risk.

An alternative probabilistic estimation of risk was also conducted using SSD curves obtained with all the species tested (see Fig. 2). This approach is not limited by the choice of AF and critical values. With that aim, data on TT values for aquatic organisms were fitted to a log-logistic model ([Van Straalen & Denneman, 1989](#)). This allowed calculations of the 5th percentile of the TT distribution (HC_5), i.e. the SMNP density affecting less than 5% of the species, and the corresponding 95% CI. If the SMNP levels do not exceed the lower end of the 95% CI of the HC_5 , then we can expect with a 95% confidence that less than 5% of the species in the community will be affected, which is considered an acceptable level in the environment ([Durán & Beiras, 2017](#)).

3 RESULTS

3.1 Exposure levels

3.1.1 Exposure levels in the river water column

The microplastic abundance of plastic particles between 10 µm and 1 mm in the water of the Thames and Elbe rivers are shown in Fig. 33 of Deliverable D2.3². This dataset was updated with previously unpublished 2023 sampling data. The main statistics of this dataset are compiled in Table 1 below. Total microplastic mass concentrations showed a high variability with no consistent seasonal trends. In 2022 levels were higher in the Thames than in the Elbe, whereas in 2023, when only hotspots were sampled, Elbe showed higher values. To produce a conservative risk assessment, the 95% percentile of the data obtained in each sampling site and the hotspot values were used to calculate the Risk Quotients.

² Statistics were recalculated using the original dataset to correct minor mistakes in the Deliverable and additional 2023 data were added.

The contents of this document are the copyright of the LABPLAS consortium and shall not be copied in whole, in part, or otherwise reproduced, used, or disclosed to any other third parties without prior written authorisation.

Table 1. Abundance of floating plastic particles between 10 µm and 1 mm in the water of the Thames and Elbe rivers.

10-1000 µm MP	Thames		Elbe		Overall
	winter	summer	winter	summer	
mean (N/m ³)	782	517	506	2716	1137
mean (µg/m ³)	922	548	1019	4804	1662
std	1876	1014	2087	9238	4520
geom mean	151	2.0	43	130	44
95% perc	3173	2142	3674	21244	15973
hotspot	Ilseworth 2022	Lechlade 2022	Dessau 2023	Cuxhaven 2023	Cuxhaven 2023
µg/m ³ in hotspot	6118	2807	6366	22041	22041

3.1.2 Exposure levels in the North Sea surface water

Table 2. Abundance of floating plastic particles between 10 µm and 1 mm in the water of the North Sea

10-1000 µm MP	North Sea		
	AL586 (winter)	AL596 (summer)	Overall
mean (N/m ³)	446	93	244
std	434	60	317
mean (µg/m ³)	302	1133	777
Std	450	1523	1194
geom mean	59	82	71
95% perc	653	2814	2451
hotspot	Station 1	Station 4	Station 4-summer
µg/m ³ in hotspot	820	3230	3230

3.1.3 Exposure levels in the river sediments

Levels of SMP in the Thames and Elbe river sediments are shown in Table 3 below. Only data in numbers were available for sediment microplastics. Assuming similar morphology of sediment and water surface microplastics, mass was estimated from the median mass/number ratio of water surface microplastics obtained in the Thames and the Elbe rivers, which was 0.343 µg per particle.

Table 3. Abundance of plastic particles between 10 µm and 1 mm in the sediments of the Thames and Elbe rivers.

10-1000 µm MP	Thames		Elbe	
	T5 London	T6 estuary	E17 Dessau	E13 estuary
mean (N/Kg)	1147	1172	3676	889
std	882	1272	2679	882
maxima (N/Kg)	2117	3362	8713	1258
maxima (µg/Kg)	726	1153	2989	431

3.1.4 Estimation of mass using actual weight records

GEOMAR kindly provided unpublished data on the mass:area ratio for a dataset of over 1000 plastic particles (mean size 2.95 ± 3.67 mm) individually measured in terms of maximum dimension, minimum dimension, area and weight. The ratio followed a log-normal distribution with an average value of 0.42 mg/mm^2 . The ratio did not show any significant trend to change with particle geometry, with a slope for the ratio vs geometry (assessed as maximum dimension/minimum dimension) close to zero (-0.016). Therefore we extrapolated this value towards the lower size range of SMP particles analysed in the LABPLAS Project and obtained mass estimations x5.2 times larger than those provided by the siMPle software.

3.1.5 Exposure levels in agricultural soils

The identification and quantification of microplastics in terrestrial environments is beyond the scope of the LABPLAS Project. However, reliable and up-to-date information is available in the scientific literature. [Büks & Kaupenjohann \(2020\)](#) recently reviewed 23 studies of microplastics in land and provided a common upper limit for agricultural sites exposed to sewage sludge and mulching film application of 4.5 mg kg^{-1} dry soil.

3.2 Toxicity thresholds

Table 4 summarizes the toxicity of the plastic litter samples tested in the LABPLAS Project. These results have been updated from those presented in the previous deliverable (D6.4).

Table 4. Summary of the toxicity tests conducted with micronized environmental plastics sampled in the different areas studied in the LABPLAS Project in Germany (GE), Spain (ES) and the United Kingdom (UK). Test species and endpoints included *Vibrio fischeri* bioluminescence, sea-urchin embryo test (SET), freshwater and marine cyanobacteria/algae population growth (*Microcystis aeruginosa* and *Skeletonema* sp. respectively), freshwater and marine invertebrate reproduction and larval survival (*Daphnia magna* and *Acartia* sp respectively), development of early life stages of freshwater and marine fish (*Danio rerio* and *Cyprinodon variegatus* respectively), soil microbiota respiration, plant seedling emergence (*Lepidium sativum*) and earthworm reproduction (*Eisenia andrei*). Results have been arbitrarily classified into categories of increasing toxicity indicated by green, yellow, orange and red. Methods have been thoroughly described in previous deliverables from D6.1 to D6.4. L: leachate test; P: particle test; n.t. not tested.

Sample			Microscale test			Habitat-specific test					
Aquatic	Month/year	ID	<i>Vibrio</i>	SET		algae	invertebrates		fish		
Kiel (GE)	May 22	093	L	L		L	L		L		
North Sea (GE)	Apr 23	125	L	L	P	L	L	P	L		
Cecebre (ES)	Jun 22	095	n.t.	L		L	L		L		
Cecebre (ES)	May 23	134	L	L	P	L	L		L	P	
Elbe (GE)	Feb 23	124	L	L	P	L	L		L	P	
Elbe (GE)	Jul 23	133	L	L	P	L	L		L	P	
Thames (UK)	Jul 22	127	L	L	P	L	L		L	P	
Thames (UK)	Jul 23	130	L	L	P	L	L		L	P	
Terrestrial	Month/year	ID	<i>Vibrio</i>	SET		soil	plants		<i>Eisenia</i>		
Abegondo (ES)	Jul 22	110	L	L		L	P	L	P	L	P
Abegondo (ES)	Mar 23	129	L	L		L	P	L	P	L	P
Abegondo (ES)	Jul 23	131	L	L		L	P	L	P	L	P

Aquatic sample ID124 (from the Elbe river banks) and terrestrial sample ID110 (from agriculture soil in Abegondo) were selected as examples of plastic litter showing some toxicity to the aquatic and terrestrial test species used, respectively.

When a concentration:response relationship was found, data were fitted to a toxicity curve and the EC₁₀ (concentration reducing by 10% the endpoint; survival, growth, reproduction) was used as an estimate of the toxicity threshold (TT, mg/L). When data did not allow curve fitting, TT was estimated as the lowest concentration significantly reducing the endpoint (LOEC). When LOEC was not identified, TT was simply delimited as above the highest level tested, and No Observed Effect Concentration (NOEC) was used for computations.

3.2.1 Aquatic species

The TT values obtained for sample ID124 were 154 mg/L for SET, 209 mg/L for copepod larvae, 220 mg/L for *Daphnia*, 431 mg/L for *Microcystis* (EC₁₀), and 490 mg/L for zebrafish (EC₁₀).

Table 5. Toxicity exerted by the most toxic freshwater plastics (ID-124) on the microscale and habitat-specific tests. L: leachate test. All leachates were made up at 10 g/L except for *Daphnia* (1 g/L). Symbols like in Tab. 4.

ID124	Microscale test		Habitat-specific test			
spp	<i>Vibrio</i>	SET	algae	<i>Acartia</i>	<i>Daphnia</i>	zebrafish
test	L	L	L	L	L	L
EC ₅₀	>10,000 (<1 TU)	625 (16 TU)	2763 (3.6 TU)	870 (11 TU)	624	700 (14.4 TU)
TT (mg/L)	>10,000	154	431	209	220	5490

3.2.2 Terrestrial species

The TT value obtained for sample ID131 was 2183 mg/L for SET, 14 mg/Kg for *Eisenia* (endpoint: number of juveniles per adult) with particles and 32 mg/Kg for *Eisenia* (endpoint: number of juveniles per adult) with leachates.

Table 6. Toxicity exerted by the most toxic terrestrial plastics (ID-131) on the microscale and terrestrial tests. L: leachate test; P: particle test.

ID131	Microscale test		Habitat-specific test					
spp	<i>Vibrio</i>	SET	Soil microbioma		Plant (<i>Lepidium sativum</i>)		<i>Eisenia andrei</i>	
test	L	L	L	P	L	P	L	P
EC ₅₀	>10,000	9804 (1.02 UT)	>1250 mg/Kg	>2250 mg/Kg	>2250 mg/L ³	>2250 mg/L	706 mg/Kg 1.77 TU	378 mg/Kg 5.75 TU
TT	>10,000	2183 (EC ₁₀)	>1250 mg/Kg	>2250 mg/Kg	>2250 mg/L	>2250 mg/L	32 mg/Kg	14 mg/Kg

3.3 Deterministic estimation of risk

As explained above, in the deterministic approach Risk Quotients (R) were calculated as

$$R = EL \times AF / TT$$

Where EL were the maximum concentrations (C) recorded in the LABPLAS Project sites studied (Tables 1 to 3). To follow a conservative approach, in this analysis we will include an AF=1000 (despite chronic effects being recorded) and will use the critical value, corresponding to the most sensitive species from each habitat.

³ LABPLAS Project Deliverable D6.4

The contents of this document are the copyright of the LABPLAS consortium and shall not be copied in whole, in part, or otherwise reproduced, used, or disclosed to any other third parties without prior written authorisation.

3.3.1 Freshwater sites

TT=220 mg/L (*Daphnia*)

Table 7. Risk Quotient (R) obtained according to the deterministic method for the freshwater sites.

10-1000 µm MP	Thames		Elbe		Overall
	winter	summer	winter	summer	
C (µg/L) 95% perc	3.173	2.142	3.674	21.244	15.973
R (AF=10)	0.0001	0.0001	0.0002	0.001	0.001
R (AF=1000)	0.014	0.010	0.017	0.097	0.073
hotspot	6.118	2.807	6.366	22.041	22.041
R (AF=10)	0.0003	0.0001	0.0003	0.001	0.001
R (AF=1000)	0.028	0.013	0.029	0.100	0.100

Maximum Risk Quotients (for AF=1000) range from 0.010 to 0.100.

If we estimate mass using the mass:area ratio rather than the siMPle software, R values would range from 0.051 to 0.521.

3.3.2 Marine sites

TT=154 mg/L (sea-urchin)

Table 8. Risk Quotient (R) obtained according to the deterministic method for the marine sites.

10-1000 µm MP	AL586 (winter)	AL596 (summer)	Overall
C (µg/m³) 95% perc	0.653	2.814	2.451
R (AF=10)	0.00004	0.0002	0.0002
R (AF=1000)	0.004	0.018	0.016
hotspot	0.820 (Station 1)	3.230 (Station 4)	3.230 (Station 4-summer)
R (AF=10)	0.00005	0.0002	0.0002
R (AF=1000)	0.005	0.021	0.021

Maximum Risk Quotients (for AF=1000) range from 0.004 to 0.021

If we estimate mass using the mass:area ratio rather than the siMPle software, R values would range from 0.021 to 0.109.

3.3.3 Sediments

Due to the limited amount of micronized environmental plastics, we performed the test with *Lumbriculus variegatus* using leachates from environmentally exposed PS (extracellular polymeric substances (EPS), 1-year

The contents of this document are the copyright of the LABPLAS consortium and shall not be copied in whole, in part, or otherwise reproduced, used, or disclosed to any other third parties without prior written authorisation.

exposure in the River Lahn conducted by BfG). Here, a 24-hour leachate induced a significantly reduced growth and reproduction at 1 g/L. Therefore, LOEC was 1 g/L (used as an estimation of TT), and NOEC was 0.5 g/L. Using the maximum SMP levels from Table 3 and AF=1000, Risk Quotients ranged from 0.431 in the Elbe estuary to 2.989 in E17-Dessau. Since these values were obtained based on a single toxicity test, they must be considered very preliminary and demand comprehensive additional research before further discussion.

3.3.4 Soil

Using the lowest TT value obtained for D131 in the *Eisenia* chronic reproduction test with particles (14 mg/Kg) and an AF=10 the Risk Quotient is already >1:

$$R = 4.5 \times 10 / 14 = 3.2 > 1$$

This means that the risk in agricultural soils fertilized with sludge and exposed to mulch film is remarkable for important organisms in soil ecology such as earthworms. This result deserves further investigation in the future to make sure plastics used in agriculture will not pose unacceptable risks to the structure of soil communities.

3.4 Probabilistic Estimation of Risk




Figure 2. SSD curves for the most toxic plastic samples in aquatic (ID124) (A), and terrestrial (ID131) (B) habitats respectively. Note that terrestrial data cannot be fit to a log-logistic model.

As shown in Fig. 2A, data on TT values for aquatic organisms fit well into a log-logistic model. This allows calculation of the 5th percentile of the TT distribution (HC₅), i.e. the SMNP density affecting less than 5% of the species, with narrow confidence intervals (CI). HC₅= 65.68 mg/L 95% CI: (21.7 – 111.5) mg/L. In contrast, as shown in Fig. 2B, data obtained for terrestrial plastics do not accurately fit this model. HC₅= 446.76 mg/L, but the 95% CI of the HC₅ estimation is unacceptably high (0.0004 - 2125) and does not allow application of this probabilistic approach. Additional ecotoxicological testing of environmental plastics with other terrestrial species beyond those used in the LABPLAS Project is needed to fill this gap.

Assuming a log-normal distribution of the SMP levels in aquatic habitats (see [Beiras & Schönemann, 2020](#)), and using the data obtained by NOC in mass units from the Great North Sea sampling (including Thames and Elbe rivers), the probability of encountering SMP levels above 21.7 mg/L (i.e. 21,700,000 µg/m³) is **R = 0.079**.

Therefore the probabilistic approach produces a R remarkably similar to the deterministic approach ($R=0.073$ for the freshwater sites and $R=0.016$ for the marine sites, using the 95% percentile and $AF=1000$), highlighting the robustness of the present assessment.

4 DISCUSSION

This deliverable provides, to the best of our knowledge, the best comprehensive effort to assess the risk of plastics in the aquatic and terrestrial environments using a combination of field studies at the European scale to assess the exposure levels, and ecotoxicological bioassays conducted with environmental plastics from the same sites to assess their toxicity thresholds. It is worth highlighting that the environmental levels recorded here include **plastic particles down to 10 μm** . This strongly contributes to shortening the frequently discussed mismatch between the size range of MP recorded in most previous studies (frequently $> 0.3 \text{ mm}$) and those reported as posing the most risk to aquatic organisms in laboratory experiments (frequently $< 20 \mu\text{m}$).

To maximize environmental protection, a number of **conservative assumptions** were adopted in the present analysis, including the following:

- We took the maximum SMP concentrations recorded in the hotspots. In fact, the 2023 sampling was conducted on the hotspots only. Therefore, this affects not only the R values calculated based on the hotspot concentrations but also those calculated based on the 95% percentiles, which for that year do not include SMP values from medium and low levels of pollution.
- In the deterministic approach, we took as TT the critical value; i.e. the value for the most sensitive species and endpoint among all the ecotoxicological bioassays conducted for each habitat.
- In the aquatic assessment, we also took the most toxic environmental plastic sample found (ID-124 from Elbe). This means we are assuming all plastics detected in WP2 sampling show the toxicity corresponding to the most toxic field sample. However, in practice, most other field plastic samples were several orders of magnitude less toxic.
- A very high AF value was also included ($AF=1000$) despite information on chronic effects was available.
- In the probabilistic approach, we used the lower end of the 95% confidence interval of the HC_5 rather than the HC_5 itself.

Even under all these extremely conservative assumptions, aquatic plastic always yields $R<1$, disregarding the approach (deterministic or probabilistic) or method used to estimate mass units from particle size.

These findings are in line with the very low-risk quotients previously obtained for coastal and oceanic ecosystems by [Beiras & Schönemann](#) (2020) and by [Adam et al.](#) (2019) in freshwaters.

In contrast, when **Risk Quotients** are calculated **for agriculture soil habitats** using the very sensitive chronic endpoint of earthworm reproduction and maximum MP levels from the literature, **unacceptably high R values** are obtained, even using a moderate AF value of 10. This deserves further attention in the regulation of plastics used for agriculture, such as mulch films, greenhouses and silos, which were among the terrestrial plastic typologies here analysed.

Eisenia's sensitivity to microplastics was about one order of magnitude higher than the toxicity thresholds for the most MP-sensitive aquatic species. Despite remarkable, this is not the main reason why R values are much higher in soil compared to both freshwater and marine environments. Most of the difference found in R between aquatic and terrestrial habitats is due to the much **higher plastic loads reported⁴ in some agricultural soils**, compared to the SMP levels recorded by the LABPLAS Project in European aquatic ecosystems.

⁴ Büks & Kaupenjohann (2020) <https://doi.org/10.5194/soil-6-649-2020>

The contents of this document are the copyright of the LABPLAS consortium and shall not be copied in whole, in part, or otherwise reproduced, used, or disclosed to any other third parties without prior written authorisation.